

# VORIZOLE

(Voriconazole for eye drops 30 mg / vial – Lyophilized Powder)

*The ultimate drug for Fungal Keratitis*



- Superior anti-fungal activity
- Lyophilized form enhances drug stability and solubility
- No side effects, established safety profile
- Economical pack
- Refrigeration not required

## DESCRIPTION

Vorizole is an antifungal drug to treat serious fungal infections. It contains 30 mg sterile lyophilized dry Voriconazole powder. To get 1% (10 mg/ml) solution, this powder has to be reconstituted with 3 ml of Water for Injection provided along with the pack.

## INDICATIONS

Vorizole is indicated for use in the treatment of fungal keratitis caused by *Aspergillus spp.*, *Fusarium spp.* and *Candida spp.* and visually devastating disease due to *Scedosporium apiospermum*

In-vitro studies have shown Voriconazole exhibits broad spectrum antifungal action against

- Moniliaceous molds - *Aspergillus fumigatus*, *Aspergillus flavus*, *Fusarium sp.*, *Paecilomyces lilacinus*, *Acremonium alabamensis*
- Dimorphic fungi - *Blastomyces dermatitidis*, *Coccidioides immitis*, *Histoplasma capsulatum*, *Penicillium marneffeii*
- Dematiaceous fungi - *Curvularia spp.*, *Scedosporium apiospermum*
- Yeast and yeast like species - *Candida albicans*, *Candida parapsilosis*, *Candida tropicalis*, *Cryptococcus neoformans*

## WARNING AND PRECAUTIONS

Vorizole is NOT FOR INJECTION. Vorizole is recommended for topical ophthalmic use only. Do not touch the nozzle tip to any surface, since this may contaminate the solution.

Voriconazole may cause visual symptoms including photophobia altered/enhanced visual perception, blurred vision and/or color vision change. The majority of visual symptoms appeared to spontaneously resolve within 60 minutes. The effect of Vorizole on visual function is not yet known if the treatment continues beyond 28 days. If treatment continues beyond 28 days, visual function including visual acuity, visual field and color perception should be monitored.

## CONTRAINDICATIONS

Vorizole is contraindicated in patients with known hypersensitivity to voriconazole or its excipients.

## ADVERSE REACTIONS

No reported adverse reaction, sometimes it may cause serious hepatic reactions. Drug interaction is a common concern.

## DOSAGE AND ADMINISTRATION

Vorizole sterile powder for eye drops should be reconstituted with 3 ml of WFI to get 10 mg / ml. (1% solution). Use the reconstituents within 7 days. After reconstitution use the drug as directed by the physician.

## SUPPLY

Supplied as 30 mg/vial in a sterile lyophilized powder form.

## STORAGE

Reconstituted solution can be stored at a temperature between 2° C and 35° C

Information published in this catalogue is subject to change without notification.